

CVS Pharmacy, Inc.

-PE Product Labels –

-continued

Drug Facts (continued)

When using this product, do not exceed recommended dosage.

Stop use and ask a doctor if:

- nervousness, dizziness, or sleeplessness occur
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- redness or swelling is present
- new symptoms occur
- fever gets worse or lasts more than 3 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed (see Overdose warning) ■ do not take more than 4 doses in 24 hours ■ swallow whole; do not crush, chew, or dissolve ■ adults and children 12 years and over: take 2 softgels with water every 4 hours ■ children under 12 years: do not use

Other information ■ store between 15-30°C (59-86°F) ■ avoid excessive heat

Inactive ingredients butylated hydroxyanisole, butylated hydroxytoluene, FD&C yellow #6, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol, sorbitol, white ink

Questions or comments? Call 1-877-763-3035 Monday-Friday 9AM-5PM EST

*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil®, NyQuil®, and LiquiCaps® are registered trademarks of The Procter and Gamble Company.

PLD-0570A F0905701

PARENTS: Learn about teen medicine abuse

www.StopMedicineAbuse.org

PLEASE RECYCLE

CVS Health.

Compare to the active ingredients in Vicks® DayQuil® and NyQuil® Cold & Flu LiquiCaps®:

Cold/Flu NDC 69842-852-24

Non-Drowsy Daytime Multi-Symptom Cold/Flu Relief

ACETAMINOPHEN Pain reliever; Fever reducer
DEXTROMETHORPHAN HBr Cough suppressant
PHENYLEPHRINE HCl Nasal decongestant

Relieves: Aches; Fever; Cough; Nasal congestion; Sore throat
Alcohol free
Antihistamine free

16 SOFTGELS**

Nighttime Multi-Symptom Cold/Flu Relief

ACETAMINOPHEN Pain reliever; Fever reducer
DEXTROMETHORPHAN HBr Cough suppressant
DOXYLAMINE SUCCINATE Antihistamine

Relieves: Aches; Fever; Cough; Sore throat; Sneezing; Runny nose

8 SOFTGELS**
24 TOTAL (Liquid-filled capsules)**

Drug Facts Nighttime Cold & Flu Softgel

Active ingredients (in each softgel)

Active ingredients (in each softgel)	Purposes
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 15 mg	Cough suppressant
Doxylamine succinate 6.25 mg	Antihistamine

Uses ■ temporarily relieves common cold and flu symptoms:
 ■ cough due to minor throat and bronchial irritation
 ■ sore throat ■ headache ■ minor aches and pains
 ■ fever ■ runny nose and sneezing

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 ■ more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin redness ■ hives ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ liver disease ■ glaucoma ■ cough that occurs with too much phlegm (mucus)
 ■ a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema
 ■ trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers

When using this product: ■ do not exceed recommended dosage ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery

Drug Facts (continued)

Stop use and ask a doctor if:

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed (see Overdose warning) ■ do not take more than 4 doses in 24 hours ■ swallow whole; do not crush, chew, or dissolve ■ adults and children 12 years and over: take 2 softgels with water every 6 hours ■ children under 12 years: do not use

Other information ■ store between 15-30°C (59-86°F) ■ avoid excessive heat

Inactive ingredients FD&C yellow #10, FD&C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol, sorbitol, white ink

Questions or comments? Call 1-877-763-3035 Monday-Friday 9AM-5PM EST

Drug Facts Daytime Cold & Flu Softgel

Active ingredients (in each softgel)

Active ingredients (in each softgel)	Purposes
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses ■ temporarily relieves common cold and flu symptoms:
 ■ cough due to minor throat and bronchial irritation
 ■ nasal congestion ■ headache
 ■ minor aches and pains ■ fever ■ sore throat

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 ■ more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin redness ■ hives ■ rash. If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ liver disease ■ heart disease ■ diabetes ■ thyroid disease ■ high blood pressure ■ cough that occurs with too much phlegm (mucus) ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Lot No.:

Exp. Date:



Compare to the active ingredients
in Theraflu® ExpressMax®
Daytime Severe Cold & Cough*

Severe Cold & Cough

NDC 59779-691-08

ACETAMINOPHEN

Pain reliever; Fever reducer

DEXTROMETHORPHAN HBr

Cough suppressant

PHENYLEPHRINE HCl

Nasal decongestant

Relief of

Nasal congestion; Cough;
Fever; Body ache;
Sore throat; Headache

DAYTIME



Berry Flavor
Alcohol content 10%
For Ages 12 & Over

8.3 FL OZ (245 mL)

P.D.-83089B
L8008861

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY
SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.**

*This product is not manufactured or distributed by GSK
Consumer Healthcare, distributor of Theraflu® ExpressMax®
Daytime Severe Cold & Cough.

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2022 CVS/pharmacy
CVS.com® 1-800-SHOP CVS V-13114

#455446



P.D.-C398B
L8008864

Drug Facts

Active ingredients (in each 30 mL)

Purposes

Acetaminophen 650 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....Cough suppressant
Phenylephrine HCl 10 mg.....Nasal decongestant

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

Uses

- temporarily relieves these symptoms due to a cold
 - minor aches and pains
 - nasal and sinus congestion
 - cough due to minor throat and bronchial irritation
- headache
- sore throat
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
 - blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are
taking the blood thinning drug warfarin.

Drug Facts (continued)

**When using this product, do not exceed
recommended dosage.**

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 6 doses (180 mL) in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device. ■ mL = milliliter
- keep dosing cup with product
- adults and children 12 years and over
 - 30 mL every 4 hours
- children under 12 years of age: do not use

Other information

- each 30 mL contains: sodium 16 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients acesulfame potassium, alcohol, citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, maltitol, propylene glycol, purified water, sodium benzoate, sodium citrate

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

PEEL CORNER FOR MORE DRUG FACTS

CVS Health. Compare to the active ingredients in Theraflu® ExpressMax® Nighttime Severe Cold & Cough*
NDC 59779-693-08

Severe Cold & Cough

ACETAMINOPHEN

Pain reliever; Fever reducer
DIPHENHYDRAMINE HCl
Antihistamine; Cough suppressant
PHENYLEPHRINE HCl
Nasal decongestant

Relief of:
Nasal congestion; Runny nose;
Cough; Fever; Body ache;
Sore throat; Headache

NIGHTTIME

Berry Flavor
Alcohol content 10%
For Ages 12 & Over

8.3 FL OZ (245 mL)

PLD-C0908
LB00885

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Theraflu® ExpressMax® Nighttime Severe Cold & Cough.

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2022 CVS/pharmacy
CVS.com® 1-800-SHOP CVS V-13114

#455443



PLD-C0908
LB00885

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg.....Pain reliever/fever reducer
Diphenhydramine HCl 25 mg.....Antihistamine/
Cough suppressant
Phenylephrine HCl 10 mg.....Nasal decongestant

Purposes

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

Uses

- temporarily relieves these symptoms due to a cold
- minor aches and pains
- headache
- sore throat
- runny nose
- sneezing
- itchy, watery eyes due to hay fever
- nasal and sinus congestion
- itching of the nose or throat
- cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
 - blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Drug Facts (continued)

Do not use ■ to make a child sleepy

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- high blood pressure
- heart disease
- diabetes
- thyroid disease
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking

- the blood thinning drug warfarin
- sedatives or tranquilizers

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

When using this product

- do not exceed recommended dosage
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- marked drowsiness may occur
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses (180 mL) in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device. ■ mL = milliliter
- keep dosing cup with product
- adults and children 12 years and over
- 30 mL every 4 hours
- children under 12 years of age: do not use

Other information

- each 30 mL contains: potassium 10 mg
- each 30 mL contains: sodium 14 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients acesulfame potassium, alcohol, citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, maltitol, propylene glycol, purified water, sodium benzoate, sodium citrate

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

2624





2626

Nighttime Cold & Flu**Drug Facts****Active ingredients (in each 30 mL)**

Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 30 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine

Purpose**Uses** temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat ■ headache ■ minor aches and pains
- fever ■ runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ hives ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease ■ glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur ■ avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. **Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ take only as directed – see Overdose warning

- only use the dose cup provided ■ do not exceed 4 doses per 24 hrs
- adults & children 12 yrs & over 30 mL every 6 hrs
- children 6 to under 12 yrs ask a doctor
- children under 4 yrs do not use

Other information

- each 30 mL contains: sodium 38 mg
- store at 20-25°C (68-77°F)

Inactive ingredients alcohol, anhydrous citric acid, D&C yellow no. 10, FD&C green no. 3, FD&C yellow no. 6, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments? 1-800-719-9260

Daytime Cold & Flu**Drug Facts****Active ingredients (in each 15 mL)**

Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Purpose**Uses**

temporarily relieves common cold/flu symptoms:

- nasal congestion ■ cough due to minor throat and bronchial irritation
- sore throat ■ headache ■ minor aches and pains ■ fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen

■ adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ hives ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease ■ heart disease ■ high blood pressure
- thyroid disease ■ diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. **Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ take only as directed – see Overdose warning

- only use the dose cup provided ■ do not exceed 4 doses per 24 hrs
- adults & children 12 yrs & over 30 mL every 4 hrs
- children 6 to under 12 yrs 15 mL every 4 hrs
- children 4 to under 6 yrs ask a doctor
- children under 4 yrs do not use

Other information

- each 15 mL contains: sodium 7 mg ■ store at 20-25°C (68-77°F)

Inactive ingredients butylated hydroxyanisole, edetate disodium, FD&C yellow no. 6, flavor, glycerin, methyl, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

Questions or comments? 1-800-719-9260



Non-Drowsy

Daytime**ACETAMINOPHEN-**

Pain reliever; Fever reducer

DEXTROMETHORPHAN HBr-

Cough suppressant

PHENYLEPHRINE HCl-

Nasal decongestant

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895

© 2022 CVS/pharmacy
CVS.com® 1-800-SHOP CVS

V-16430

100% money back
guaranteed.
CVS.com/returnpolicy

*These products are not manufactured or distributed
by Pfecta & Gemma, distributor of Vicks® DayQuil®
Cold & Flu and Vicks® NyQuil® Cold & Flu.

CVS Health. Compare to the active ingredients in Vicks® DayQuil® Cold & Flu®

Non-Drowsy Daytime

NDC 59779-656-40

ACETAMINOPHEN-
Pain reliever; Fever reducer
DEXTROMETHORPHAN HBr-
Cough suppressant
PHENYLEPHRINE HCl-
Nasal decongestant

Relieves:
Aches, Fever,
Sore throat, Cough,
Nasal congestion

COLD & FLU

**ALCOHOL &
ANTIHISTAMINE FREE**

12 FL OZ (355 mL)

: 65640 17 FL

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2022 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-16430

100% money back guaranteed.
CVS.com/returnpolicy

Empty & Replace Cap

PLASTIC BOTTLE PLASTIC CUP

0 50428 32557 5

Drug Facts

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses ■ temporarily relieves common cold/flu symptoms:
■ cough due to minor throat and bronchial irritation
■ nasal congestion ■ minor aches and pains
■ sore throat ■ headache ■ fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if ■ adult takes more than 4,000 mg of acetaminophen in 24 hours ■ child takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening
■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

PEEL BACK AT CORNER FOR MORE INFORMATION

: 65640 17 81

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

ADHESIVE AREA
■ NO VARNISH ■ NO TYPE

Drug Facts (continued)



Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults) ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ take only as directed - see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs

ADHESIVE AREA
■ NO VARNISH ■ NO TYPE





2631



Compare to the active ingredients in
Vicks® DayQuil® Severe Cold & Flu*

Severe

MAXIMUM STRENGTH

Daytime Softgels

Severe Cold/Flu Relief

ACETAMINOPHEN - Pain reliever, Fever reducer

DEXTROMETHORPHAN HBr - Cough suppressant

GUAIFENESIN - Expectorant

PHENYLEPHRINE HCl - Nasal decongestant

Relieves:

Headache, Fever, Sore throat, Minor aches
& pains, Cough, Chest congestion, Nasal/Sinus
congestion & Sinus pressure

Non-drowsy

Alcohol free

Antihistamine free



Actual Size

24 SOFTGELS





Compare to the
active ingredients in
Vicks® Day-Quil®
Severe Cold & Flu*



2632 Compare to the
active ingredients in
Vicks® NyQuil®
Severe Cold & Flu*

Non-Drowsy

Severe Daytime

COLD & FLU RELIEF

ACETAMINOPHEN

Pain reliever/Fever reducer

DEXTROMETHORPHAN HBr

Cough suppressant

GUAIFENESIN

Expectorant

PHENYLEPHRINE HCl

Nasal decongestant

MAXIMUM STRENGTH

Relieves:

Aches; Fever; Cough; Nasal
congestion; Sore throat;
Chest congestion

Alcohol free
Antihistamine
free



Actual Size

12 SOFTGELS

Severe Nighttime

COLD & FLU RELIEF

ACETAMINOPHEN

Pain reliever/Fever reducer

DEXTROMETHORPHAN HBr

Cough suppressant

DOXYLAMINE SUCCINATE

Antihistamine

PHENYLEPHRINE HCl

Nasal decongestant

MAXIMUM STRENGTH

Relieves:

Aches; Fever; Sore throat;
Cough; Sneezing; Runny
nose; Nasal/Sinus
congestion &
Sinus pressure



Actual Size

12 SOFTGELS 24 TOTAL

[illegible]



Actual Size

CVS Health

Effervescent Cold Relief

ASPIRIN (NSAID)
Pain reliever; Fever reducer

CHLORPHENIRAMINE MALEATE
Antihistamine

PHENYLEPHRINE BITARTRATE
Nasal decongestant

Original Flavor

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you are allergic to aspirin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you are taking sedatives or tranquilizers
- you have glaucoma
- diabetes
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- been placed on a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- presently taking a prescription drug
- taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

When using this product do not take more than 8 tablets (adults and children 12 years and over) in a 24-hour period or as directed by doctor.

- do not use more than directed
- you may get drowsy
- avoid alcoholic drinks
- excitability may occur, especially in children
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs
- nervousness, dizziness or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely

Drug Facts (continued)

directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not use more than directed (see overdose warning)
- adults and children 12 years and over: take 2 tablets completely dissolved in 4 oz of water every 4-6 hours
- do not take more than 8 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

each tablet contains:

- sodium 464 mg
- phenylephrine bitartrate 9 mg per tablet

Store at room temperature (59°-86°F)

Inactive ingredients

acesulfame potassium, aspartame, citric acid, docusate sodium, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

This product is not manufactured, distributed or licensed by the maker of Alka-Seltzer Plus® Cold.

DO NOT USE IF PRINTED PACKETS ARE TORN OR PUNCTURED

VALUE SIZE

CVS Health

Compare to the active ingredients in Alka-Seltzer Plus® Cold Relief!

Original Flavor

NDC 69842-768-36

Effervescent Cold Relief

ASPIRIN (NSAID)
Pain reliever; Fever reducer

CHLORPHENIRAMINE MALEATE
Antihistamine

PHENYLEPHRINE BITARTRATE
Nasal decongestant

Fast relief of:

- Nasal congestion; Runny nose; Sneezing; Headache;
- Body ache; Sore throat;
- Sinus pain & pressure



Actual Size Tablet on Top Panel

36 TABLETS

Drug Facts

Active ingredients (in each effervescent tablet)

Aspirin 325 mg (NSAID)
Chlorpheniramine maleate 2 mg
Phenylephrine bitartrate 7.5 mg

Purpose

Pain reliever; fever reducer
Antihistamine
Nasal decongestant
Nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves these symptoms of the common cold:
 - fever
 - sneezing
 - nasal congestion
 - sore throat
 - headaches
 - minor aches and pains
 - runny nose
 - sinus congestion and pressure

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: hives, facial swelling, asthma (wheezing), shock

Stomach bleeding warning: This product contains a non-steroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinning (anticoagulant) or steroid drug, take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed.

PN: 87368 CH C1
REV: 02/17

Distributed by:
CVS Pharmacy, Inc.
One CVS Drive
Woonsocket, RI 02895
© 2017 CVS/pharmacy
CVS.com®
1-800-SHOP CVS
V-17432

CVS Quality
Money Back Guarantee

#182129



0 50428 58107 0

CVS Health Compare to active ingredients in Theraflu® Daytime Severe Cold & Cough*

Daytime Relief

NDC 59779-926-08

Flu & Severe Cold

ACETAMINOPHEN
Pain reliever; Fever reducer

DEXTROMETHORPHAN HBr
Cough suppressant

PHENYLEPHRINE HCl
Nasal decongestant

Relief of:
Nasal congestion;
Cough; Fever; Body
ache; Sore throat;
Headache



Cherry Flavor
Alcohol content 10%
For Ages 12 & Over

8.3 FL OZ (245 mL)

50428-37797

PLD-E264B LB002803

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by Novartis Consumer Health, Inc., distributor of Theraflu® Daytime Severe Cold & Cough.

Failure to follow the drug facts warnings could result in serious consequences.

#455446



Distributed by: CVS Pharmacy, Inc.

One CVS Drive
Woonsocket, RI 02895
© 2015 CVS/pharmacy
CVS.com® 1-800-SHOP CVS

CVS Quality
Money Back Guarantee

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....Cough suppressant
Phenylephrine HCl 10 mg.....Nasal decongestant

Purposes

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

Uses

- temporarily relieves these symptoms due to a cold
- minor aches and pains ■ headache
- nasal congestion ■ sore throat
- cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure

Drug Facts (continued)

whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease ■ heart disease ■ thyroid disease
- diabetes ■ high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device. ■ mL = milliliter
- keep dosing cup with product
- adults and children 12 years and over
 - 30 mL every 4 hours
- children under 12 years of age: do not use

Other information

- each 30 mL contains: sodium 16 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Drug Facts (continued)

Inactive ingredients acesulfame potassium, alcohol, citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, maltitol, propylene glycol, purified water, sodium benzoate, sodium citrate

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Drug Facts

Active ingredients (in each softgel)

Active ingredients (in each softgel)	Purposes
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 6 mg	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - headache
 - nasal congestion
 - sore throat
 - cough
 - minor aches and pains
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- diabetes
- high blood pressure
- heart disease
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- a persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warnings: Taking more than the recommended dose (eventual) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 softgels in any 24-hour period
- adults and children 12 years of age and older: take 2 softgels every 4 hours
- children under 12 years of age: do not use

Other information

- store between 20-25°C (68-77°F)
- avoid excessive heat
- swallow whole; do not crush, chew, or dissolve

Inactive ingredients: FD&C red #40, FD&C yellow #6, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, titanium dioxide

Questions or comments?
Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

(This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Cold, Flu & Sore Throat.)

TAMPER EVIDENT DO NOT USE IF BUBBLER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

CVS Health.

ACETAMINOPHEN - Pain reliever, Fever reducer
DEXTROMETHORPHAN HBr - Cough suppressant
GUAIFENESIN - Expectorant
PHENYLEPHRINE HCl - Nasal decongestant

Compare to the active ingredients in Maximum Strength Mucinex® Fast-Max® Cold, Flu & Sore Throat†

NDC 69842-969-16

MAXIMUM STRENGTH

Cold, Flu & Sore Throat

ACETAMINOPHEN - 325 mg - Pain reliever; Fever reducer
DEXTROMETHORPHAN HBr - 10 mg - Cough suppressant
GUAIFENESIN - 200 mg - Expectorant
PHENYLEPHRINE HCl - 5 mg - Nasal decongestant

Relieves:

- Cough
- Nasal & chest congestion
- Headache & fever

Thins & loosens mucus

For Ages 12 Years & Over
Alcohol Free



Actual Size

16 SOFTGELS**
(**LiquiL-filled capsules)

CVS Health.

MAXIMUM STRENGTH

Cold, Flu & Sore Throat

#177088



0 50428 54767 0

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2020 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-16/25

CVS Quality
Money Back Guarantee

PLD-ES89A
FD008159

Lot No:
Exp. Date:

Nighttime Cold & Flu (continued)

Drug Facts (continued)
Ask a doctor or pharmacist before use if you are:

- taking the blood thinning drug warfarin
- using sedatives or tranquilizers

When using this product:

- do not use more than directed
- excitability may occur, especially in children
- musical drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if:

- nausea or vomiting is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions

- do not take more than 12 softgels (Daytime and Nighttime) in any 24-hour period
- adults and children 12 years of age and older: take 2 softgels every 4 hours
- children under 12 years of age: do not use
- when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information

- swallow whole; do not crush, chew, or dissolve
- store between 15-30°C (59-86°F)
- avoid excessive heat

Inactive ingredients D&C yellow #10, FD&C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, polyethylene glycol, purified water, citric acid, sodium hydroxide, sorbitol, sorbitol, titanium dioxide
 *contains one or more of these ingredients
Questions or comments?
 Call 1-877-753-3835 Monday-Friday 9AM-5PM EST

Daytime Severe Cold

Drug Facts
Active ingredients (in each softgel)
 Acetaminophen 325 mg.....Pain reliever/fever reducer
 Dextromethorphan HBr 10 mg.....Cough suppressant
 Pseudoephedrine HCl 5 mg.....Expectorant
 Phenylephrine HCl 5 mg.....Nasal decongestant
Purposes

- temporarily relieves these common cold and flu symptoms:
 - headache
 - runny nose
 - cough
 - minor aches and pains
 - helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of both mucus and make cough more productive
 - temporarily reduces fever

Uses

- temporarily relieves these common cold and flu symptoms:
 - headache
 - runny nose
 - cough
 - minor aches and pains
 - helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of both mucus and make cough more productive
 - temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Severe throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have:

- liver disease
- diabetes
- high blood pressure
- heart disease
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- a persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Daytime Severe Cold (continued)

Drug Facts (continued)
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.
When using this product, do not use more than directed.
Stop use and ask a doctor if:

- nausea, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 softgels (Daytime and Nighttime) in any 24-hour period
- adults and children 12 years of age and older: take 2 softgels every 4 hours
- children under 12 years of age: do not use
- when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information

- swallow whole; do not crush, chew, or dissolve
- store between 15-30°C (59-86°F)
- avoid excessive heat

Inactive ingredients FD&C red #40, FD&C yellow #6, gelatin, glycerin, mannitol, polyethylene glycol, polyethylene glycol, purified water, sorbitol, sorbitol, titanium dioxide
 *may contain this ingredient
Questions or comments?
 Call 1-877-753-3835 Monday-Friday 9AM-5PM EST
 *This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Day Severe Cold and Maximum Strength Mucinex® Fast-Max® Night Cold & Flu.

DAY & NIGHT TWIN PACK

CVS Health.
MAXIMUM STRENGTH Daytime Severe Cold
MAXIMUM STRENGTH Nighttime Cold & Flu
ACETAMINOPHEN - 325 mg Pain reliever; Fever reducer
DEXTROMETHORPHAN HBr - 10 mg Cough suppressant
GUAFENESIN - 200 mg Expectorant
PHENYLEPHRINE HCl - 5 mg Nasal decongestant
Relieves aches, fever & sore throat
Controls cough
Relieves nasal & chest congestion
Thins & loosens mucus
For Ages 12 Years & Older
Alcohol Free
 Actual Size
16 SOFTGELS**

MAXIMUM STRENGTH Nighttime Cold & Flu
ACETAMINOPHEN - 325 mg Pain reliever; Fever reducer
DEXTROMETHORPHAN HBr - 10 mg Cough suppressant
DOXYLAMINE SUCCINATE - 6.25 mg Antihistamine
PHENYLEPHRINE HCl - 5 mg Nasal decongestant
Relieves aches, fever & sore throat
Controls cough
Relieves nasal congestion
Relieves runny nose & sneezing
For Ages 12 Years & Older
Alcohol Free
 Actual Size
8 SOFTGELS**
24 TOTAL (**Liquid-filled capsules)

#177184

0 50428 54778 6
PARENTS: P.D. 9421C F305487
www.CVSHealth.com

TAMPER EVIDENT: DO NOT USE IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.
KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.
Distributed by: CVS Pharmacy, Inc.
 One CVS Drive, Woonsocket, RI 02895
 © 2019 CVS/pharmacy
 CVS.com® 1-800-SHOP CVS V-16425
CVS Quality
 Every Week Guarantee
Lot No.:
Exp. Date:

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING



Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2020 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-36611
Money. Back Guarantee

Drug Facts (continued)

■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin
When using this product do not use more than directed
Stop use and ask a doctor if
■ nervousness, dizziness, or sleeplessness occur
■ pain, nasal congestion, or cough gets worse or lasts more than 7 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
■ new symptoms occur
■ cough comes back or occurs with rash or headache that lasts
These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ **do not take more than directed (see Overdose warning)**
■ do not take more than 12 caplets in any 24-hour period
■ adults and children 12 years of age and over: take 2 caplets every 4 hours
■ children under 12 years of age: do not use

Other information

■ store between 20-25°C (68-77°F) in a dry place
■ retain outer package for complete product information and warnings

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, crospovidone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

Drug Facts

Active ingredients (in each caplet) **Purpose**
Acetaminophen 325 mg.....Pain reliever
Dextromethorphan HBr 10 mg.....Cough suppressant
Guaifenesin 200 mg.....Expectorant
Phenylephrine HCl 5 mg.....Nasal decongestant
Uses ■ temporarily relieves:
■ headache
■ nasal congestion
■ minor aches and pains
■ sinus congestion and pressure
■ cough
■ promotes nasal and/or sinus drainage
■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
■ more than 12 caplets in 24 hours, which is the maximum daily amount
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks daily while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening
■ blisters
■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

■ diabetes
■ heart disease
■ high blood pressure
■ thyroid disease
■ trouble urinating due to an enlarged prostate gland
■ cough that occurs with too much phlegm (mucus)

MAXIMUM STRENGTH

Mucus Relief Sinus Pressure, Pain & Cough



Compare to the active ingredients in Mucinex® Sinus-Max® Pressure, Pain & Cough*

MAXIMUM STRENGTH

NDC 69842-796-02

Mucus Relief Sinus Pressure, Pain & Cough

ACETAMINOPHEN - Pain reliever
DEXTROMETHORPHAN HBr - Cough suppressant
GUAIFENESIN - Expectorant
PHENYLEPHRINE HCl - Nasal decongestant

Relieves:

- Headache
- Sinus pressure & congestion
- Cough

Thins & loosens mucus

20 CAPLETS

Ages 12 Years & Over

Actual Size



F119002CVS_RO

#938776



Drug Facts (continued)
Questions or comments?
1-844-705-4384

*This product is not manufactured or distributed by Hechtel Benckiser, distributor of Mucinex® Sinus-Max® Pressure, Pain & Cough.
Maximum Strength Per 4-Hour Dose

CVS Quality
Money Back Guarantee

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2020 CVS Pharmacy
CVS.com® 1-800-SHOP CVS
V-36611

**TAMPER EVIDENT: DO NOT USE IF
PACKAGE IS OPENED OR IF BLISTER
UNIT IS TORN, BROKEN OR SHOWS
ANY SIGNS OF TAMPERING**

MAXIMUM STRENGTH

Mucus Relief Sinus Severe Congestion & Pain



Compare to the active ingredients
in Mucinex® Sinus-Max®
Severe Congestion & Pain*

NDC 69842-786-02

MAXIMUM STRENGTH

Mucus Relief Sinus Severe Congestion & Pain

ACETAMINOPHEN - Pain reliever
GUAIFENESIN - Expectorant
PHENYLEPHRINE HCl - Nasal decongestant

Relieves:

- Sinus congestion
 - Headache
- Thins & loosens mucus

20 CAPLETS

Ages 12 Years & Over

Actual Size



Drug Facts	
Active ingredients (in each caplet) Acetaminophen 325 mg Pain reliever Guaifenesin 200 mg Expectorant Phenylephrine HCl 5 mg Nasal decongestant	Purpose Pain reliever Expectorant Nasal decongestant
Uses temporarily relieves: ■ nasal congestion ■ sinus congestion and pressure ■ promotes nasal and/or sinus drainage ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive	Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 12 caplets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks daily while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.
Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	Ask a doctor before use if you have ■ liver disease ■ high blood pressure ■ trouble urinating due to an enlarged prostate gland ■ cough that occurs with too much phlegm (mucus)
Directions ■ do not take more than directed (see Overdose warning) ■ do not take more than 12 caplets in any 24-hour period ■ adults and children 12 years and older: take 2 caplets every 4 hours ■ children under 12 years of age: do not use	Other information ■ store between 20-25°C (68-77°F) in a dry place ■ retain outer package for complete product information
Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, croscopolone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide	

*This product is not manufactured or distributed by Mucinex® Sinus-Max® distributor of Mucinex® Sinus-Max®.
Severe Congestion & Pain.
Maximum Strength Per 4-Hour Dose

Drug Facts (continued)
Questions or comments?
1-844-705-4384

F116702CVS_R0

#938777



0 50428 33235 1

Multi-Symptom Cold Drug Facts (cont.)

Drug Facts (continued)

Other information

- each 5 mL contains: sodium 4 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, disodium EDTA, FD&C red #40, flavor, glycerin, propyl galate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

*This product is not manufactured or distributed by Rockitt Benckiser, distributor of Children's Mucinex® Multi-Symptom Cold.

Multi-Symptom Nighttime Cold Drug Facts (cont.)

Drug Facts (continued)

- with any other drug containing diphenhydramine even one used on the skin
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product
- to make a child sleepy

Ask a doctor before use if your child has

- liver disease ■ heart disease ■ high blood pressure
- diabetes ■ thyroid disease ■ glaucoma
- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if your child is

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days
- redness or swelling is present
- new symptoms occur
- fever gets worse or lasts more than 3 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Drowsiness warning)
- do not give more than 5 doses in any 24-hour period
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 days unless directed by a doctor
- measure only with dosing cup provided. Do not use any other dosing device. ■ mL = milliliter
- dose as follows or as directed by a doctor
- children 6 to under 12 years of age: 10 mL every 4 hours
- children under 6 years of age: do not use

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Drowsiness warning)
- do not give more than 5 doses in any 24-hour period
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 days unless directed by a doctor
- measure only with dosing cup provided. Do not use any other dosing device. ■ mL = milliliter
- dose as follows or as directed by a doctor
- children 6 to under 12 years of age: 10 mL every 4 hours
- children under 6 years of age: do not use

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Drowsiness warning)
- do not give more than 5 doses in any 24-hour period
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 days unless directed by a doctor
- measure only with dosing cup provided. Do not use any other dosing device. ■ mL = milliliter
- dose as follows or as directed by a doctor
- children 6 to under 12 years of age: 10 mL every 4 hours
- children under 6 years of age: do not use

Multi-Symptom Nighttime Cold Drug Facts

Drug Facts

Active ingredients (in each 10 mL) Purposes

Acetaminophen 325 mg Pain reliever/fever reducer
Diphenhydramine HCl 12.5 mg Antihistamine/Cough suppressant
Phenylephrine HCl 5 mg Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms
- minor aches and pains ■ headache ■ sore throat
- sneezing ■ runny nose
- nasal congestion ■ cough
- controls cough to help your child get to sleep
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash

Severe throat warning: If your throat is sore, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Multi-Symptom Nighttime Cold Drug Facts (cont.)

Drug Facts (continued)

Other information

- each 10 mL contains: sodium 5 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl galate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?
Call 1-877-755-3855 Monday-Friday 9AM-5PM EST

*This product is not manufactured or distributed by Rockitt Benckiser, distributor of Children's Mucinex® Night Time Multi-Symptom Cold.

DAY & NIGHT TWIN PACK

CVS Health.

Compare to the active ingredients in Children's Mucinex® Night Time Multi-Symptom Cold*
NDC 69842-820-08

Children's Multi-Symptom Cold

DEXTROMETHORPHAN HBr 5 mg
Cough suppressant
GUAIFENESIN 100 mg
Expectorant
PHENYLEPHRINE HCl 2.5 mg
Nasal decongestant

Relieves:
Nasal & chest congestion
Soothes cough
Thins & loosens mucus

Children's Multi-Symptom Nighttime Cold

ACETAMINOPHEN 325 mg
Pain reliever/Fever reducer
DIPHENHYDRAMINE HCl 12.5 mg
Antihistamine/Cough suppressant
PHENYLEPHRINE HCl 5 mg
Nasal decongestant

Relieves:
Nasal & chest congestion
Reduces fever
Soothes cough & sore throat

Berry Flavor Liquid
Ages 4 to 11 Years
Dosage cup included

Berry Flavor Liquid
Ages 6 to 11 Years
Dosage cup included

TWO 4 FL OZ (118 mL) BOTTLES TOTAL - 8 FL OZ (236 mL)

Multi-Symptom Cold Drug Facts

Drug Facts

Active ingredients (in each 5 mL) Purposes

Dextromethorphan HBr 5 mg Cough suppressant
Guaifenesin 100 mg Expectorant
Phenylephrine HCl 2.5 mg Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of both mucus and mucus made more productive
- temporarily relieves
- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the intensity of coughing
- the impulse to cough to help your child get to sleep
- nasal congestion due to a cold
- stuffy nose

Warnings

Do not use

- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if your child has

- heart disease ■ high blood pressure
- diabetes ■ thyroid disease
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

When using this product, do not use more than directed.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough lasts more than 7 days, comes back, or occurs with a fever, rash or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than 6 doses in any 24-hour period
- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- keep dosing cup with product
- dose as follows or as directed by a doctor ■ mL = milliliter

Age	Dose
children 6 years to under 12 years	10 mL every 4 hours
children 4 years to under 6 years	5 mL every 4 hours
children under 4 years	do not use

#177011



0 50428 54247 7

PLD-83518
PLD-8350C FC004176Lot No.:
Exp. Date:**PARENTS:**
Read and understand this information
www.StayMedSafe.com

Drug Facts (continued)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days
- redness or swelling is present
- new symptoms occur
- fever gets worse or lasts more than 3 days
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions ■ this product does not contain directions or complete warnings for adult use

- do not give more than directed (see Overdose warning)
- do not give more than 5 doses in any 24-hour period
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 doses unless directed by a doctor
- measure only with dosing cup provided. Do not use any other dosing device.
- mL = milliliter
- dose as follows or as directed by a doctor
- children 6 to under 12 years of age: 10 mL every 4 hours
- children under 6 years of age: do not use

Other information ■ each 10 mL contains: sodium 5 mg

- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients: citric acid, dextromethorphan HBr, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?
Call 1-877-753-2936 Monday-Friday 9AM-5PM EST

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Children's Mucinex® Night Time Multi-Symptom Cold.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2018 CVS/pharmacy
CVS.com® 1-800-SHOP CVS V-13114

CVS Quality
Money Back Guarantee



Package Contains One Bottle

Actual Size

CVS Health

Compare to the active ingredients in Children's Mucinex® Night Time Multi-Symptom Cold*

NDC 69842-506-04

Children's Multi-Symptom Nighttime Cold

ACETAMINOPHEN 325 mg
Pain reliever/Fever reducer

DIPHENHYDRAMINE HCl 12.5 mg
Antihistamine/Cough suppressant

PHENYLEPHRINE HCl 5 mg
Nasal decongestant

Relieves:
Nasal congestion

**Reduces fever
Soothes cough & sore throat**

Ages 6 to 11 Years



Mixed Berry Flavor
Dosage cup included

Actual Bottle Size on Side Panel

4 FL OZ (118 mL)

Drug Facts

Active ingredients (in each 10 mL)	Purposes
Acetaminophen 325 mg	Pain reliever/fever reducer
Diphenhydramine HCl 12.5 mg	Antihistamine/Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms
- minor aches and pains
- headache
- sore throat
- sneezing
- runny nose
- nasal congestion
- cough
- controls cough to help your child get to sleep
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ hives ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- with any other drug containing diphenhydramine even one used on the skin
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product
- to make a child sleepy

Ask a doctor before use if your child has

- liver disease
- heart disease
- high blood pressure
- diabetes
- thyroid disease
- glaucoma
- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if your child is

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product ■ do not use more than directed

- drowsiness may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

#194343



PJL-K032C FC030007

Lot No:

Exp. Date:



Children's Multi-Symptom Cold & Fever



Children's Multi-Symptom Cold & Fever

ACETAMINOPHEN
Pain reliever/Fever reducer

DEXTROMETHORPHAN HBr
Cough suppressant

GUAIFENESIN
Expectorant

PHENYLEPHRINE HCl
Nasal decongestant
ORAL SOLUTION

Relieves:

Minor aches & pains, Fever, Cough,
Nasal congestion, Chest congestion,
Stuffy nose, Thins & loosens mucus

Alcohol free



Mixed Berry Flavored Liquid
Ages 6 to 11 Years

4 FL OZ (118 mL)

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY
SEAL UNDER CAP IS BROKEN OR MISSING

Package Contains One Bottle

Actual Size

Drug Facts

KEEP OUTER PACKAGE FOR
COMPLETE PRODUCT INFORMATION

Active ingredients (in each 10 mL)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves common cold and flu symptoms:
 - minor aches and pains
 - sore throat
 - headache
 - stuffy nose
 - the intensity of coughing
 - nasal congestion
 - cough due to minor throat and bronchial irritation
 - the impulse to cough to help your child get to sleep
 - temporarily reduces fever
 - helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount

- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if the child has

- liver disease
- high blood pressure
- heart disease
- thyroid disease
- diabetes
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

Drug Facts (continued)

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed
- this product does not contain directions or complete warnings for adult use
- do not give more than 5 doses in any 24-hour period
- mL = milliliter
- only use the dose cup provided
- repeat dose every 4 hours while symptoms last
- dose as follows or as directed by a doctor
- children 6 to under 12 years: 10 mL in dosing cup provided
- children under 6 years: do not use

Other information

- each 10 mL contains: sodium 5 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients anhydrous citric acid, FD&C blue #1, FD&C red #40, flavors, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sorbitol, sucralose

Questions or comments? 1-800-426-9391

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2023 CVS/pharmacy

CVS.com®
1-800-SHOP CVS
V-19849



50844 REV0123A07536

B-0231-075-36-H
REV0123A07536

#895140



No print/No varnish
Lot & Exp date

2643

CVS Health

Nasal Decongestant PE

PHENYLEPHRINE HCl TABLETS
Nasal decongestant

MAXIMUM STRENGTH

Relieves:

- Sinus pressure
- Nasal congestion

72 TABLETS 10 mg EACH

Non-Drowsy

Compare to the active ingredient in Sudafed PE® Sinus Congestion*

Actual Size

Drug Facts

Active ingredient (in each tablet) Phenylephrine HCl 10 mg Nasal decongestant

Purpose Relieves nasal congestion due to the common cold.

Uses

- Temporarily relieves sinus congestion and pressure
- May relieve or other upper respiratory allergies

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- diabetes
- thyroid disease
- high blood pressure

When using this product do not exceed recommended dosage.

- difficulty in urination due to enlargement of the prostate gland
- nervousness, dizziness, or sleeplessness occur

Stop use and ask a doctor if

- symptoms do not improve within 7 days or occur with fever

Questions or comments? 1-800-426-9391

Inactive ingredients croscarmellose sodium, dextrose monohydrate, dibasic calcium phosphate dihydrate, FD&C red #40, lecithin, magnesium stearate, malodextrin, microcrystalline cellulose, silicon dioxide, sodium carboxymethylcellulose, sodium citrate dihydrate, titanium dioxide

Other information

- TAMPER EVIDENT: DO NOT USE IF OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Directions

- adults and children 12 years and over: take 1 tablet every 4 hours. Do not take more than 6 tablets in 24 hours.
- children under 12 years: ask a doctor

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2021 CVS/pharmacy
CVS.com® 1-800-SHOP CVS V-19849

B-0231-453-23-H
REV0820A45323

NO PRINT / NO VARNISH AREA
LOT NO. & EXP. DATE
331010
0 50428 127285 5

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not give more than 6 doses in 24 hours unless directed by a doctor
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product ■ mL = milliliter
- shake well before using

Age	Dose
children 6 to under 12 years of age	10 mL every 4 hours
children 4 to under 6 years of age	do not use unless directed by a doctor
children under 4 years of age	do not use

Other information ■ each 5 mL contains: sodium 3 mg

■ store between 20-25°C (68-77°F). Do not refrigerate. ■ protect from light

Inactive ingredients: acesulfame potassium, citric acid, EDTA disodium, FD&C blue #1, FD&C red #40, flavor, maltitol, propylene glycol, purified water, sodium benzoate, sodium citrate

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2019 CVS/pharmacy
CVS.com® 1-800-SHOP CVS V-13114



PLD-B320E LB005843



Children's Nighttime Cold & Cough

NDC 69842-379-04

DIPHENHYDRAMINE HCl
Antihistamine; Cough suppressant
PHENYLEPHRINE HCl
Nasal decongestant

Relieves:

- Cough
- Runny, stuffy nose
- Itchy throat

Alcohol free

For Ages 6 to 11 Years Grape Flavor



4 FL OZ (118 mL)

Drug Facts

Active ingredients (in each 5 mL)

Diphenhydramine HCl 6.25 mg

Phenylephrine HCl 2.5 mg

Purposes

Antihistamine; Cough suppressant

Nasal decongestant

Uses ■ temporarily relieves:

- sneezing ■ runny nose ■ itchy nose or throat
- itchy, watery eyes due to hay fever ■ nasal and sinus congestion
- cough due to minor throat and bronchial irritations as may occur with a cold

Warnings

Do not use ■ in a child under 4 years of age

■ in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

■ with any other product containing diphenhydramine, even one used on skin

■ for the purpose of making your child sleepy

Ask a doctor before use if the child has ■ heart disease

■ high blood pressure ■ thyroid disease ■ diabetes ■ glaucoma

■ cough that occurs with too much phlegm (mucus)

■ chronic cough that lasts, or as occurs with asthma

■ a breathing problem such as chronic bronchitis

Ask a doctor or pharmacist before use if the child is taking sedatives or tranquilizers.

When using this product ■ do not exceed recommended dosage

■ marked drowsiness may occur ■ excitability may occur, especially in children

■ sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if ■ nervousness, dizziness, or sleeplessness occurs

■ symptoms do not improve within 7 days or occur with a fever

■ cough persists for more than 7 days, comes back, or occurs with a fever, rash, or persistent headache. These could be signs of a serious condition.

CVS Health. Compare to the active ingredients in Theraflu® ExpressMax® Nighttime Severe Cold & Cough*

Severe Cold & Cough
ACETAMINOPHEN
Pain reliever; Fever reducer
DIPHENHYDRAMINE HCl
Antihistamine; Cough suppressant
PHENYLEPHRINE HCl
Nasal decongestant

NIGHTTIME

Berry Flavor
Alcohol content 10%
For Ages 12 & Over

8.3 FL OZ (245 mL)

NDC 59779-693-08

P.D.-C399B
L8008885

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Theraflu® ExpressMax® Nighttime Severe Cold & Cough.

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2022 CVS/pharmacy
CVS.com® 1-800-SHOP CVS V-13114

#455443

P.D.-C399B
L8008885

0 50428 25333 5

Drug Facts**Active ingredients (in each 30 mL)**

Acetaminophen 650 mg.....Pain reliever/fever reducer
Diphenhydramine HCl 25 mg.....Antihistamine/
Cough suppressant
Phenylephrine HCl 10 mg.....Nasal decongestant

Purposes**PEEL CORNER FOR MORE DRUG FACTS****Drug Facts (continued)****Uses**

- temporarily relieves these symptoms due to a cold
 - minor aches and pains ■ headache
 - sore throat ■ runny nose ■ sneezing
 - itchy, watery eyes due to hay fever
 - nasal and sinus congestion
 - itching of the nose or throat
 - cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Drug Facts (continued)

Do not use ■ to make a child sleepy

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease ■ glaucoma
- high blood pressure ■ heart disease
- diabetes ■ thyroid disease
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking

- the blood thinning drug warfarin
- sedatives or tranquilizers

PEEL CORNER FOR MORE DRUG FACTS**Drug Facts (continued)****When using this product**

- do not exceed recommended dosage
- alcohol, sedatives, and tranquilizers may increase drowsiness ■ avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery ■ marked drowsiness may occur
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, cough, or nasal congestion gets worse or lasts more than 7 days ■ new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)**Directions**

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses (180 mL) in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device. ■ mL = milliliter
- keep dosing cup with product
- adults and children 12 years and over
 - 30 mL every 4 hours
- children under 12 years of age: do not use

Other information

- each 30 mL contains: potassium 10 mg
- each 30 mL contains: sodium 14 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients acesulfame potassium, alcohol, citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, maltitol, propylene glycol, purified water, sodium benzoate, sodium citrate

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST



Compare to the
active ingredients in
Vicks® NyQuil® Severe*
NDC 69842-800-40

MAXIMUM STRENGTH

Nighttime Severe

ACETAMINOPHEN-
Pain reliever; Fever reducer
DEXTROMETHORPHAN HBr-
Cough suppressant
DOXYLAMINE SUCCINATE-
Antihistamine
PHENYLEPHRINE HCl-
Nasal decongestant

Relieves:

Aches, Fever, Sore throat,
Cough, Sneezing, Runny nose,
Nasal & Sinus congestion,
Sinus pressure

COLD & FLU

Original Flavor
ALCOHOL 10%

12 FL OZ (355 mL)

: 60940 17 F3

PARENTS:

Learn about teen medicine abuse
www.StopMedicineAbuse.org

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2022 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-16430

100% money back
guaranteed.
CVS.com/returnpolicy

Empty &
Replace Cap



PLASTIC
BOTTLE

PLASTIC
CUP

how2recycle.info

Drug Facts

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg.....	Pain reliever/ fever reducer
Dextromethorphan HBr 20 mg.....	Cough suppressant
Doxylamine succinate 12.5 mg.....	Antihistamine
Phenylephrine HCl 10 mg.....	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms: ■ sinus congestion and pressure ■ nasal congestion ■ minor aches and pains ■ headache ■ runny nose and sneezing ■ sore throat ■ cough to help you sleep ■ fever ■ cough due to minor throat and bronchial irritation ■ reduces swelling of nasal passages ■ promotes nasal and/or sinus drainage ■ temporarily restores freer breathing through the nose

Warnings

Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening

■ blisters ■ rash
If a skin reaction occurs,
stop use and seek medical
help right away.

**PEEL BACK AT
CORNER FOR MORE
INFORMATION**

: 60940 17 B1

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
■ liver disease ■ heart disease

Drug Facts (continued)

When using this product ■ do not use more than directed ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. Overdose warnings:
In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ take only as directed - see Overdose warning ■ only use the dose cup

RESERVE AREA
NO VARNISH • NO TYPE

ADHESIVE AREA
NO VARNISH • NO TYPE

2647

AD
NO COATING

- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

NO COATING

provided ■ do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information ■ each 30 mL contains: sodium 44 mg ■ store at 20-25°C (68-77°F)

Inactive ingredients alcohol, anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C green #3, FD&C yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose

Questions?
1-800-719-9260

*This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® NyQuil® Severe.

NO COATING

Non-Drowsy
Severe Daytime
COLD & FLU RELIEF

Compare to the active ingredients in Vicks VapoGel™ Severe Cold & Flu*

Non-Drowsy
Severe Nighttime
COLD & FLU RELIEF

Compare to the active ingredients in Vicks VapoGel™ Severe Cold & Flu*

Severe Daytime
COLD & FLU RELIEF

Non-Drowsy

ACETAMINOPHEN
Pain reliever/fever reducer
DEXTROMETHORPHAN HBr
Cough suppressant
GUAFENESIN
Expectorant
PHENYLEPHRINE HCl
Nasal decongestant

Relieves:
Aches; Fever; Cough; Nasal congestion; Sore throat; Chest congestion
Alcohol free
Antihistamine free

MAXIMUM STRENGTH

Actual Size

12 SOFTGELS

Severe Nighttime
COLD & FLU RELIEF

Non-Drowsy

ACETAMINOPHEN
Pain reliever/fever reducer
DEXTROMETHORPHAN HBr
Cough suppressant
DOXYLAMINE SUCCINATE
Antihistamine
PHENYLEPHRINE HCl
Nasal decongestant

Relieves:
Aches; Fever; Sore throat; Cough; Sneezing; Runny nose; Nasal/Sinus congestion & Sinus pressure

MAXIMUM STRENGTH

Actual Size

12 SOFTGELS 24 TOTAL

THIS PRODUCT IS PACKAGED IN A CHILD-RESISTANT AND TAMPER-EVIDENT PACKAGE. USE ONLY IF BLISTERS ARE INTACT.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

DO NOT TAKE THESE PRODUCTS AT THE SAME TIME

NIGHTTIME SEVERE COLD & FLU

Drug Facts

Active ingredients (in each softgel)	Purposes
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Doxylamine succinate 6.25 mg.....	Antihistamine
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses

- temporarily relieves common cold/flu symptoms:
- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- minor aches & pains
- headache
- fever
- sore throat
- runny nose & sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

Warnings

Liver warning This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 softgels in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

DAYTIME SEVERE COLD & FLU

Drug Facts

Active ingredients (in each softgel)	Purposes
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Guafenesin 200 mg.....	Expectorant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses

- temporarily relieves common cold/flu symptoms:
- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- minor aches & pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Liver warning This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 softgels in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

NIGHTTIME SEVERE COLD & FLU

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen ask a doctor or pharmacist.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleep

Ask a doctor before use if you have

DAYTIME SEVERE COLD & FLU

Drug Facts (continued)

Allergy alert Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen ask a doctor or pharmacist.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional

LOT:
EXP:



0 50428 36769 8

#970808

<p>CVS Quality  Made in India V-85249 Distributed by: CVS Pharmacy, Inc. One CVS Drive, Woonsocket, RI 02895 © 2020 CVS/Pharmacy CVS.com 1-800-SHOP CVS</p>	<p>Questions or comments? Call toll free: 1-888-333-9792</p>
<p>Drug Facts (continued) This product is not manufactured or distributed by Vicks® DayQuil® and NyQuil® Severe Cold & Flu. Procter & Gamble, owner of the registered trademarks</p>	<p>Drug Facts (continued) purified water, sorbitol sorbitan solution, titanium dioxide</p>
<p>Other information ■ store at room temperature</p> <p>Inactive ingredients D&C Yellow #10, FD&C Blue #1, gelatin, polyethylene glycol 400, povidone K80, propylene glycol, purified water, shellac, sorbitol sorbitan, sodium hydroxide, titanium dioxide</p>	<p>Other information ■ store at room temperature</p> <p>Inactive ingredients FD&C Red #40, FD&C Yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide</p>
<p>Directions ■ take only as directed - see Overdose warning ■ do not exceed 8 softgels per 24 hrs adults & children 12 years & over 2 softgels with water every 4 hours children 4 to under 12 years ask a doctor children under 4 years do not use</p> <p>Drug Facts (continued) Stop use and ask a doctor if: ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p>	<p>Directions ■ take only as directed - see Overdose warning ■ do not exceed 8 softgels per 24 hrs adults & children 12 years & over 2 softgels with water every 4 hours children 4 to under 12 years ask a doctor children under 4 years do not use</p> <p>Drug Facts (continued) Stop use and ask a doctor if: ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p>
<p>NIGHTTIME SEVERE COLD & FLU</p>	<p>DAYTIME SEVERE COLD & FLU</p>

C050428367698



R 3090

Severe Allergy & Sinus Headache

MAXIMUM STRENGTH

Caplets

CVS Health

Severe Allergy & Sinus Headache

MAXIMUM STRENGTH

Caplets

CVS Health

B-0231-543-09-H
REV0918C54309

Drug Facts

KEEP OUTER PACKAGE FOR
COMPLETE PRODUCT INFORMATION

Active ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever
Diphenhydramine HCl 25 mg	Antihistamine
Phenylephrine HCl 5 mg	Nasal decongestant

Uses ■ temporarily relieves these symptoms of hay fever or other upper respiratory allergies: ■ headache
■ nasal congestion ■ sneezing ■ runny nose
■ sinus congestion and pressure ■ itchy, watery eyes
■ itching of the nose or throat ■ minor aches and pains

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ with any other product containing diphenhydramine, even one used on skin
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ if you have ever had an allergic reaction to this product or any of its ingredients

Drug Facts (continued)

Ask a doctor before use if you have
■ glaucoma ■ heart disease ■ high blood pressure
■ difficulty in urination due to enlargement of the prostate gland
■ a breathing problem such as emphysema or chronic bronchitis ■ liver disease ■ thyroid disease ■ diabetes

Ask a doctor or pharmacist before use if you are

■ taking the blood thinning drug warfarin
■ taking sedatives or tranquilizers
When using this product
■ do not exceed recommended dosage
■ excitability may occur, especially in children
■ marked drowsiness may occur
■ alcohol, sedatives, and tranquilizers may increase drowsiness
■ use caution when driving a motor vehicle or operating machinery ■ avoid alcoholic beverages

Stop use and ask a doctor if

■ nervousness, dizziness, or sleeplessness occur
■ pain or nasal congestion gets worse or lasts more than 7 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present ■ new symptoms occur

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ do not take more than directed
■ adults and children 12 years and over
■ take 2 caplets every 4 hours
■ do not take more than 10 caplets in 24 hours
■ children under 12 years: ask a doctor

Other information

■ see end flap for expiration date and lot number

Drug Facts (continued)

■ **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients corn starch, croscarmellose

Drug Facts (continued)

sodium, crospovidone, FD & C blue #1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, povidone, silicon dioxide, stearic acid, titanium dioxide, triacetin

Questions or comments? 1-800-426-9391



#343025-PE

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
OPENED OR IF BLISTER UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS OF TAMPERING

50844

REV0918C54309

no print / no varnish area
- - - - - for use & exp. date - - - - -

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2018 CVS/pharmacy
CVS.com • 1-800-SHOP CVS
V-18815

CVS Quality
Money Back Guarantee







Do not use if printed neckband is broken or missing.

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02805
© 2019 CVS/pharmacy
CVS.com® 1-800-SHOP CVS

V-16430



Drug Facts

Active ingredients (in each 20 mL)	Purpose
Acetaminophen 650 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....	Cough suppressant
Guaifenesin 400 mg.....	Expectorant
Phenylephrine HCl 10 mg.....	Nasal decongestant

Uses ■ temporarily relieves these common cold and flu symptoms:
■ cough ■ nasal congestion ■ minor aches and pains ■ sore throat ■ headache ■ stuffy nose ■ sinus congestion and pressure
■ temporarily reduces fever ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters

■ rash ■ If a skin reaction occurs, stop use and seek medical help right away.

PEEL CORNER TO READ COMPLETE
DRUG FACTS AND INFORMATION



: 7W63D 17 B3

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have ■ liver disease ■ heart disease ■ diabetes ■ high blood pressure ■ thyroid disease ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if ■ nervousness, dizziness or sleeplessness occur ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed (see Overdose warning) ■ do not take more than 5 doses in any 24-hour period ■ measure only with dosing cup provided ■ do not use dosing cup with other products ■ dose as follows or as directed by a doctor ■ adults and children 12 years of age and older: 20 mL, in dosing cup provided every 4 hours ■ children under 12 years of age: do not use

Other information

■ each 20 mL contains: sodium 7 mg
■ store at 20-25°C (68-77°F)
■ do not refrigerate

Inactive ingredients anhydrous citric acid, benzyl alcohol, edetate disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, triethyl citrate, xanthan gum

Questions or comments?
1-800-719-9260

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Mucinex® Fast-Max® Severe Cold.



Compare to the active ingredients in Theraflu® Multi-Symptom Severe Cold

Severe

NDC 69642-949-91

Multi-Symptom Severe Cold

ACETAMINOPHEN
Pain reliever; Fever reducer
DEXTROMETHORPHAN HBr
Cough suppressant
PHENYLEPHRINE HCl
Nasal decongestant

Relieves:

- Nasal congestion
- Sore throat pain
- Cough
- Headache
- Body ache
- Fever



Green Tea & Honey Lemon Flavors

6 PACKETS



Severe

Multi-Symptom Severe Cold

CVS Health™ Multi-Symptom Severe Cold provides complete relief of your severe cold symptoms in a soothing hot liquid formula.

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

CONVENIENT RESEALING TAB

DO NOT USE IF PRINTED PACKETS ARE TORN OR PUNCTURED

Drug Facts

Active ingredients (in each packet)

Active ingredients (in each packet)	Purposes
Acetaminophen 500 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold:
 - minor aches and pains
 - minor sore throat pain
 - headache
 - nasal and sinus congestion
 - cough due to minor throat and bronchial irritation
 - temporarily reduce fever

Warnings

Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 2 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- in a child under 12 years of age
- if you have ever had an allergic reaction to this product or any of its ingredients
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- do not use more than directed (see overdose warning)
- take every 4 hours, while symptoms persist. Do not take more than 6 packets in 24 hours unless directed by a doctor.

Age	Dose
adults and children 12 years of age and over	one packet
children under 12 years of age	do not use

Other information

- each packet contains: potassium 10 mg and sodium 25 mg
- phenylephrine HCl contains phenylephrine 22 mg per packet
- store at 20-25°C (68-77°F). Protect product from heat and moisture.

Inactive ingredients acsulfate potassium, anhydrous citric acid, aspartame, colloidal silicon dioxide, D&G yellow #10, FD&C blue #1, FD&C red #40, flavors, maltodextrin, pregelatinized starch, sodium citrate, sucrose, tribasic calcium phosphate

Questions or comments?
1-800-718-4280

*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Theraflu® Multi-Symptom Severe Cold.

Distributed by:
CVS Pharmacy, Inc.
One CVS Drive
Woonsocket, RI 02895
© 2018 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
Made in Mexico V-16430

CVS Quality
Money Back Guarantee

OPEN OTHER END
#633103



LOT NO.

EXP.

81494 37 CL

No print/No varnish
- Lot & Exp date -

No Print/No Varnish
Lot and Expiration No.

8 CAPLETS

Drug Facts	
Active ingredients (in each softgel)	Purposes
Acetaminophen 325 mg.....Pain reliever/fever reducer Dextromethorphan HBr 10 mg.....Cough suppressant Guaifenesin 200 mg.....Expectorant Phenylephrine HCl 5 mg.....Nasal decongestant	
Uses	
<ul style="list-style-type: none"> temporarily relieves these common cold and flu symptoms: <ul style="list-style-type: none"> headache nasal congestion sore throat cough minor aches and pains helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive temporarily reduces fever 	
Warnings	
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: <ul style="list-style-type: none"> more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product 	
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> skin reddening blisters rash If a skin reaction occurs, stop use and seek medical help right away.	
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	
Do not use	
<ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. 	
Ask a doctor before use if you have	
<ul style="list-style-type: none"> liver disease diabetes high blood pressure heart disease thyroid disease trouble urinating due to an enlarged prostate gland a persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema cough that occurs with too much phlegm (mucus) 	
Drug Facts (continued)	
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.	
When using this product, do not use more than directed.	
Stop use and ask a doctor if	
<ul style="list-style-type: none"> nervousness, dizziness, or sleeplessness occur pain, nasal congestion, or cough gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. 	
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children.	
Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
Directions	
do not take more than directed (see Overdose warning) <ul style="list-style-type: none"> do not take more than 12 softgels in any 24-hour period adults and children 12 years of age and older: take 2 softgels every 4 hours children under 12 years of age: do not use 	
Other information	
<ul style="list-style-type: none"> store between 20-25°C (68-77°F) avoid excessive heat swallow whole; do not crush, chew, or dissolve 	
Inactive ingredients	
FD&C red #40, FD&C yellow #6, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, titanium dioxide	
Questions or comments?	
Call 1-877-753-3835 Monday-Friday 9AM-5PM EST	
*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Severe Cold Liquid Gels.	
TAMPER EVIDENT: DO NOT USE IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.	
KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.	

Severe Cold

MAXIMUM STRENGTH

CVS Health



Compare to the active ingredients
in Maximum Strength Mucinex®
Fast-Max® Severe Cold Liquid Gels**
NDC 69842-905-16

MAXIMUM STRENGTH Severe Cold

Acetaminophen - Pain reliever, Fever reducer
Dextromethorphan HBr - Cough suppressant
Guaifenesin - Expectorant
Phenylephrine HCl
Nasal decongestant

MULTI-SYMPTOM

Controls cough

Thins & loosens mucus

Relieves nasal & chest congestion

Relieves aches, fever & sore throat



FOR AGES 12+
Alcohol Free

Actual Size

16 SOFTGELS[†] (†Liquid-filled capsules)

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2020 CVS/pharmacy
CVS.com®
1-800-SHOP CVS V-16425



PARENTS:
Learn about teen medicine risks
www.StopMedicineAbuse.org



Lot No.:
Exp. Date:

Drug Facts	
Active ingredients (in each softgel)	Purposes
Acetaminophen 325 mg.....Pain reliever/fever reducer Dextromethorphan HBr 10 mg.....Cough suppressant Guaifenesin 200 mg.....Expectorant Phenylephrine HCl 5 mg.....Nasal decongestant	
Uses	
<ul style="list-style-type: none"> temporarily relieves these common cold and flu symptoms: <ul style="list-style-type: none"> headache nasal congestion sore throat cough minor aches and pains helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive temporarily reduces fever 	
Warnings	
<p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:</p> <ul style="list-style-type: none"> more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> skin redness hives rash <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use</p> <ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> liver disease diabetes high blood pressure heart disease thyroid disease trouble urinating due to an enlarged prostate gland a persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema cough that occurs with too much phlegm (mucus) 	
Drug Facts (continued)	
<p>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.</p> <p>When using this product, do not use more than directed.</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> nervousness, dizziness, or sleeplessness occur pain, nasal congestion, or cough gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children.</p> <p>Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p>	
Directions	
<ul style="list-style-type: none"> do not take more than directed (see Overdose warning) do not take more than 12 softgels in any 24-hour period adults and children 12 years of age and older: take 2 softgels every 4 hours children under 12 years of age: do not use 	
Other information	
<ul style="list-style-type: none"> store between 15-30°C (59-86°F) avoid excessive heat swallow whole; do not crush, chew, or dissolve 	
Inactive ingredients	FD&C red #40, FD&C yellow #6, gelatin, glycerin, mannitol*, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol, sorbitol, titanium dioxide
*may contain this ingredient	
Questions or comments?	Call 1-877-753-3935 Monday-Friday 9AM-5PM EST
<p>This product is not manufactured or distributed by Beckitt. Beckitt, distributor of Maximum Strength Mucinex® Fast-Max® Day Severe Cold.</p> <p>TAMPER EVIDENT: DO NOT USE IF BLISTER UNIT IS TORN, SPOKEN OR SHOWS ANY SIGNS OF TAMPERING.</p> <p>KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.</p>	

CVS Health.

Severe Cold

MAXIMUM STRENGTH

Compare to the active ingredients in Maximum Strength Mucinex® Fast-Max® Day Severe Cold†
NDC 69842-837-16

Severe Cold

ACETAMINOPHEN - Pain reliever, Fever reducer
DEXTROMETHORPHAN HBr - Cough suppressant
GUAIFENESIN - Expectorant
PHENYLEPHRINE HCl - Nasal decongestant

MAXIMUM STRENGTH

Multi-Symptom

- Relieves aches, fever & sore throat
- Controls cough
- Relieves nasal & chest congestion
- Thins & loosens mucus

For Ages 12 Years & Older
Alcohol Free



Actual Size

16 SOFTGELS**
(**Liquid-filled capsules)

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2019 CVS/pharmacy
CVS.com®
1-800-SHOP CVS V-16425



PARENTS:
Learn about your medicine here
www.StopMedicineAbuse.org



Lot No.:
Exp. Date:

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredients (in each caplet)

Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses ■ temporarily relieves:

- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the intensity of coughing
- nasal congestion due to a cold
- the impulse to cough to help you get to sleep
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- high blood pressure
- heart disease
- diabetes
- difficulty in urination due to enlargement of the prostate gland
- thyroid disease
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough persists more than 7 days, tends to recur, or is accompanied by a fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not take more than directed

- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years and over: take 2 caplets every 4 hours
- children under 12 years: do not use

Other information ■ each caplet contains: sodium 3 mg

- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

how2recycle.info

PAPER BOX

MULTILAYER TRAY

*This product is not manufactured or distributed by RB Health (US) LLC, owner of the registered trademark Maximum Strength Mucinex® FAST-MAX® Severe Congestion & Cough. 150% more tablets than Maximum Strength Mucinex® FAST-MAX® Severe Congestion & Cough, 20 count package.

Drug Facts (continued)

Inactive ingredients corn starch, FD&C blue #2 aluminum lake, FD&C red #40 aluminum lake, magnesium stearate, malcododrin, microcrystalline cellulose, polyethylene glycol polysorbate 80, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments? 1-800-428-9391

CVS Health.

Multi-Symptom

Severe Congestion & Cough

50% MORE TABLETS THAN THE NATIONAL BRAND*

CVS Health.

Compare to the active ingredients in Maximum Strength Mucinex® FAST-MAX® Severe Congestion & Cough*

Multi-Symptom

Severe Congestion & Cough

DEXTROMETHORPHAN HBr
Cough suppressant
GUAIFENESIN - Expectorant
PHENYLEPHRINE HCl
Nasal decongestant

MAXIMUM STRENGTH

Relieves:
Nasal & chest congestion
Controls cough
Thins & loosens mucus

For Ages 12+


Actual Size

30 COATED CAPLETS

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Distributed by:
CVS Pharmacy, Inc.
One CVS Drive
Woonsocket, RI 02895
© 2023 CVS/pharmacy
CVS.com®
1-800-SHOP CVS
V-19849

100% money back guaranteed.
CVS.com/returnpolicy

B-0231-648-01-H
REV0423D64801


#972449

FPO 80%
UPC# 050428393789

X XXXXXX XXXXXX X

No print/No varnish Area
Lot & Expiry date

50844 REV0423D64801



Drug Facts (continued)

Warnings (Nighttime only)

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: blisters, rash, skin redness.

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. (Nighttime only)
- if you have ever had an allergic reaction to this product or any of its ingredients (Nighttime only)
- with any other product containing diphenhydramine, even one used on skin (Nighttime only)

Ask a doctor before use if you have

- heart disease
- diabetes
- thyroid disease
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- high blood pressure

Drug Facts (continued)

- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis (Nighttime only)
- glaucoma (Nighttime only)
- liver disease (Nighttime only)

Ask a doctor or pharmacist before use if you are (Nighttime only)

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not exceed recommended dosage
- excitability may occur, especially in children (Nighttime only)
- alcohol, sedatives, and tranquilizers may increase drowsiness (Nighttime only)
- use caution when driving a motor vehicle or operating machinery (Nighttime only)
- marked drowsiness may occur (Nighttime only)

Stop use and ask a doctor if:

- nausea, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever (Daytime only)
- itchiness or swelling is present (Nighttime only)
- pain, nasal congestion, or cough gets worse or lasts more than 7 days (Nighttime only)
- new symptoms occur (Nighttime only)
- fever gets worse or lasts more than 3 days (Nighttime only)
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Drug Facts (continued)

Directions

Do not take DAYTIME and NIGHTTIME products at the same time.

- do not take more than directed
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years and over: take 2 caplets every 4 hours
- children under 12 years: do not use

Other information

- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients (Daytime only) corn starch, FD&C blue #2 aluminum lake, FD&C red #40 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyisobutyl alcohol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide.

Inactive ingredients (Nighttime only) corn starch, croscarmellose sodium, croscopolone, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, iron oxide yellow, magnesium stearate, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium bicarbonate, stearic acid, talc, titanium dioxide.

Drug Facts (continued)

Questions or comments? 1-800-426-9391

Distributed by: CVS Pharmacy, Inc.
 One CVS Drive, Woonsocket, RI 02895
 © 2022 CVS/pharmacy
 CVS.com® 1-800-SHOP CVS
 V-19849

CVS Quality
 Money Back Guarantee

Do not take the Daytime and Nighttime caplets at the same time.

50844
 REV1021B64869401
 R-0231-648694-01DF









Compare to the active ingredients in
Theraflu® Nighttime Severe Cold & Cough™

Severe





Compare to the active ingredients in
Theraflu® Nighttime Severe Cold & Cough™

CONVENIENT RESEALING TAB

NDC 58779-760-91

NIGHTTIME Severe Cough & Cold

ACETAMINOPHEN
Pain reliever; Fever reducer
DIPHENHYDRAMINE HCl
Antihistamine; Cough suppressant
PHENYLEPHRINE HCl
Nasal decongestant

Relieves:

- Nasal congestion
- Cough
- Runny nose
- Sneezing
- Body ache
- Sore throat pain
- Headache / fever



Honey Lemon
Infused with
White Tea Flavors

6 PACKETS

NIGHTTIME Severe Cough & Cold

CVS Health™ Severe Cough & Cold Nighttime provides complete relief of your cough and severe cold symptoms. This soothing hot liquid formula is alcohol free and tastes great.

For daytime relief, try
Daytime Severe Cough & Cold.

*This product is not manufactured or distributed by Novartis Consumer Health, Inc., distributor of Theraflu® Nighttime Severe Cold & Cough.

DO NOT USE IF PRINTED PACKETS ARE TORN OR PUNCTURED

Drug Facts

Active ingredients (in each packet)

Acetaminophen 650 mg	Pain reliever/fever reducer
Diphenhydramine HCl 25 mg	Antihistamine/cough suppressant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold:
 - minor aches and pains
 - minor sore throat pain
 - headache
 - nasal and sinus congestion
 - runny nose
 - sneezing
 - itchy nose or throat
 - itchy, watery eyes due to hay fever
 - cough
 - cough soothes minor throat and bronchial irritation
 - temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Severe throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- in a child under 12 years of age
- if you have ever had an allergic reaction to this product or any of its ingredients
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- high blood pressure
- thyroid disease
- liver disease
- diabetes
- heart disease
- glaucoma

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not exceed recommended dosage
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occurs
- redness or swelling is present
- fever gets worse or lasts more than 3 days
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Purposes

Pain reliever/fever reducer

Antihistamine/cough suppressant

Nasal decongestant

Directions

- do not use more than directed (see overdose warning)
- take every 4 hours, while symptoms persist. Do not take more than 5 packets in 24 hours unless directed by a doctor.

Age	Dose
adults and children 12 years of age and over	one packet
children under 12 years of age	do not use

Other information

- each packet contains: potassium 10 mg and sodium 25 mg
- phenylephrine HCl: contains phenylephrine 10 mg per packet
- store at 20-25°C (68-77°F). Protect product from heat and moisture.

Inactive ingredients acetaminophen, anhydrous citric acid, aspartame, colloidal silicon dioxide, D&C yellow #10, FD&C blue #1, FD&C red #40, flavors, methylcellulose, pregelatinized starch, sodium citrate, sucrose, tribasic calcium phosphate

Questions or comments?
1-800-778-5285

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2017 CVS Pharmacy
CVS.com 1-800-SHOP-CVS
Made in Mexico
V-16430


OPEN OTHER END



CODE
AREA

96491 17 C4

2662



Compare to the active ingredients
in Mucinex® Fast-Max® Severe
Congestion & Cough*

Severe Cough & Congestion


DEXTROMETHORPHAN HBr - Cough suppressant
GUAIFENESIN - Expectorant
PHENYLEPHRINE HCl - Nasal decongestant

MAXIMUM STRENGTH

Multi-Symptom

- Relieves nasal & chest congestion
- Controls cough
- Thins & loosens mucus

For Ages 12 & Over
Alcohol Free



6 FL OZ (177 mL)

F-004-45
REV B

PARENTS:
Learn about our products and services
www.ShopMyCVSStore.org

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2023 CVS/pharmacy CVS.com®
1-800-SHOP CVS V-19849

#895137

FPO 80%
UPC# 050428403044

X XXXXXXX XXXXXX X

Drug Facts TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Active ingredients (in each 20 mL)	Purpose
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing

No Print / No Varnish Area
Lot # and Exp. Info

Drug Facts (continued)

- the impulse to cough to help you get to sleep
- nasal congestion due to a cold

Warnings
Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- diabetes
- high blood pressure
- thyroid disease
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough persists more than 7 days, tends to recur, or is accompanied by a fever, rash, or persistent headache. These could be signs of a serious condition.

HINGE

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- do not take more than 6 doses in any 24-hour period
- mL = milliliter; FL OZ = fluid ounce
- only use the dose cup provided
- adults and children 12 years and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years: do not use

Other information




- each 20 mL contains: sodium 10 mg
- use by expiration date on package
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients anhydrous citric acid, FD&C blue #1, FD&C red #40, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sorbitol, sucralose, xanthan gum

Questions or comments?
1-800-426-9391

*This product is not manufactured or distributed by RB Health (US) LLC, owner of the registered trademark Mucinex® Fast-Max® Severe Congestion & Cough. 50844 REV0121B00445

HINGE



Do not use if printed neckband is broken or missing
*This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® NyQuil® Severe.

Distributed by: **CVS Pharmacy, Inc.**
One CVS Drive, Woonsocket, RI 02895
© 2018 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-16430

CVS Quality
Money Back Guarantee

0 50428125913 9 #954407

: 76334 17 F5

CVS Health

Compare to the active ingredients in Vicks® NyQuil® Severe*

Severe

NDC 59779-763-34

Nighttime

COLD & FLU RELIEF

ACETAMINOPHEN - Pain reliever; Fever reducer
DEXTROMETHORPHAN HBr - Cough suppressant
DOXYLAMINE SUCCINATE - Antihistamine
PHENYLEPHRINE HCl - Nasal decongestant

MAXIMUM STRENGTH

Relieves:
Aches; Fever; Sore throat;
Cough; Sneezing; Runny nose;
Nasal/sinus congestion
& sinus pressure

Berry Flavor
Alcohol free

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

8 FL OZ (237 mL)

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....	Cough suppressant
Doxylamine succinate 12.5 mg.....	Antihistamine
Phenylephrine HCl 10 mg.....	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms:

■ sinus congestion and pressure	■ nasal congestion
■ minor aches and pains	■ headache
■ runny nose and sneezing	■ sore throat
■ cough to help you sleep	■ fever
■ cough due to minor throat and bronchial irritation	
■ reduces swelling of nasal passages	
■ promotes nasal and/or sinus drainage	
■ temporarily restores freer breathing through the nose	

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

PEEL BACK AT CORNER FOR MORE INFORMATION

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen ("prescription or nonprescription"). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease ■ heart disease ■ high blood pressure
- thyroid disease ■ diabetes ■ glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin

When using this product ■ **do not use more than directed**

- excitability may occur, especially in children
- marked drowsiness may occur ■ avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Drug Facts (continued)

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present ■ new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ take only as directed – see Overdose warning

- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information ■ each 30 mL contains: sodium 41 mg
■ store at 20-25°C (68-77°F)

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions or comments? 1-800-719-9260

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

These products are not manufactured or distributed by Procter & Gamble, distributor of Vicks® DayQuil® Severe and Vicks® NyQuil® Severe.

TWIN PACK

CVS Health™

Compare to the active ingredients
in Vicks® DayQuil® Severe*

Severe

Non-Drowsy

Daytime

COLD & FLU RELIEF

ACETAMINOPHEN - Pain reliever; Fever reducer
DEXTROMETHORPHAN HBr - Cough suppressant
GUAIFENESIN - Expectorant
PHENYLEPHRINE HCl - Nasal decongestant

MAXIMUM STRENGTH

Relieves:
Aches; Fever; Cough;
Nasal congestion;
Sore throat;
Chest congestion



Alcohol free
Antihistamine free

TWO 12 FL OZ (355 mL) UNITS

CVS Health™

Compare to the active ingredients
in Vicks® NyQuil® Severe*

Severe

NDC 59779-597-02

Nighttime

COLD & FLU RELIEF

ACETAMINOPHEN - Pain reliever; Fever reducer
DEXTROMETHORPHAN HBr - Cough suppressant
DOXYLAMINE SUCCINATE - Antihistamine
PHENYLEPHRINE HCl - Nasal decongestant

MAXIMUM STRENGTH

Relieves:
Aches; Fever; Sore throat;
Cough; Sneezing;
Runny nose; Nasal/sinus
congestion &
Sinus pressure





Berry Flavor
Alcohol free

TOTAL 24 FL OZ (1.5 PT) (710 mL)

CVS Quality
Money Back Guarantee

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2018 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-16430

DO NOT USE IF PRINTED NECKBAND IS
BROKEN OR MISSING

Drug Facts (continued)

Inactive ingredients: butylated hydroxyanisole, edetate disodium, FD&C yellow #6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum.

Questions or comments? 1-800-719-9260

Drug Facts (continued)

Inactive ingredients: antihydroxychloride, citric acid, edetate disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum.

Questions or comments? 1-800-719-9260

Drug Facts	Nighttime Severe Cold & Flu	Drug Facts	Daytime Severe Cold & Flu
Active ingredients (in each 30 mL)		Active ingredients (in each 15 mL)	
Purpose		Purpose	
Acetaminophen 650 mg.....Pain reliever/fever reducer		Acetaminophen 325 mg.....Pain reliever/fever reducer	
Dextromethorphan HBr 20 mg.....Cough suppressant		Dextromethorphan HBr 10 mg.....Cough suppressant	
Doxylamine succinate 12.5 mg.....Antihistamine		Guaifenesin 200 mg.....Expectorant	
Phenylephrine HCl 10 mg.....Nasal decongestant		Phenylephrine HCl 5 mg.....Nasal decongestant	
Uses temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ sinus congestion and pressure ■ minor aches and pains ■ headache ■ fever ■ sore throat ■ runny nose and sneezing ■ cough due to minor throat and bronchial irritation ■ cough to help you sleep ■ reduces swelling of nasal passages ■ promotes nasal and/or sinus drainage ■ temporarily restores freer breathing through the nose		Uses temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ sinus congestion and pressure ■ cough due to minor throat and bronchial irritation ■ minor aches and pains ■ headache ■ fever ■ sore throat ■ reduces swelling of nasal passages ■ temporarily restores freer breathing through the nose ■ promotes nasal and/or sinus drainage ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive	
Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.		Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if ■ adult takes more than 4,000 mg of acetaminophen in 24 hours ■ child takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product. Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	
Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ If you have ever had an allergic reaction to this product or any of its ingredients.		Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ If you have ever had an allergic reaction to this product or any of its ingredients.	
Ask a doctor before use if you have ■ liver disease ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem such as emphysema or chronic bronchitis ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ a sodium-restricted diet		Ask a doctor before use if you have ■ liver disease ■ heart disease ■ thyroid disease ■ diabetes ■ high blood pressure ■ trouble urinating due to an enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema	
Ask a doctor or pharmacist before use if you are ■ taking sedatives or tranquilizers ■ taking the blood-thinning drug warfarin		Ask a doctor or pharmacist before use if you are ■ taking the blood-thinning drug warfarin	
When using this product ■ do not use more than directed ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness		When using this product ■ do not use more than directed	
Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.		Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults) ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.	
If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.		If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
Directions ■ take only as directed - see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs		Directions ■ take only as directed - see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs	
adults & children 12 yrs & over 30 mL every 4 hrs		adults & children 12 yrs & over 30 mL every 4 hrs	
children 4 to under 12 yrs ask a doctor		children 6 to under 12 yrs 15 mL every 4 hrs	
children under 4 yrs do not use		children 4 to under 6 yrs ask a doctor	
children under 4 yrs do not use		children under 4 yrs do not use	
Other information ■ each 30 mL contains: sodium 41 mg ■ store at 20-25°C (68-77°F)		Other information ■ each 15 mL contains: sodium 6 mg ■ store at 20-25°C (68-77°F). Do not refrigerate. ►	

h3 27 20245 :



Children's Cold & Cough Daytime

Drug Facts

Active ingredients (in each 5 mL) **Purposes**
Dextromethorphan HBr 5 mg.....Cough suppressant
Phenylephrine HCl 2.5 mg.....Nasal decongestant

Uses

- temporarily relieves
- nasal and sinus congestion
- cough due to minor throat and bronchial irritation as may occur with a cold

Warnings

- Do not use
- In a child under 4 years of age
- In a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough that occurs with too much phlegm (mucus)
- chronic cough that lasts, or as occurs with asthma

When using this product, do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occurs
- symptoms do not improve within 7 days or occur with fever
- cough persists more than 7 days, comes back or occurs with a fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not give more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- shake well before using

Age	Dose
children 6 to under 12 years of age	10 mL every 4 hours
children 4 to under 6 years of age	5 mL every 4 hours
children under 4 years of age	do not use

Children's Cold & Cough Daytime (continued)

Drug Facts (continued)

Other information

- each 5 mL contains: sodium 2 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients acesulfame potassium, alcohol, benzoic acid, citric acid, disodium EDTA, FD&C red #40, flavor, maltitol, propylene glycol, purified water, sodium citrate

Children's Nighttime Cold & Cough

Drug Facts

Active ingredients **Purposes**
Diphenhydramine HCl 6.25 mg.....Antihistamine/Cough suppressant
Phenylephrine HCl 2.5 mg.....Nasal decongestant

Uses

- temporarily relieves
- sneezing
- runny nose
- itchy nose or throat
- itchy, watery eyes due to hay fever
- nasal and sinus congestion
- cough due to minor throat and bronchial irritations as may occur with a cold

Warnings

- Do not use
- In a child under 4 years of age
- In a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- with any other product containing diphenhydramine, even one used on skin
- for the purpose of making your child sleepy

Ask a doctor before use if the child has

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- chronic cough that lasts, or as occurs with asthma
- a breathing problem such as chronic bronchitis

Ask a doctor or pharmacist before use if the child is taking sedatives or tranquilizers.

Children's Nighttime Cold & Cough (continued)

Drug Facts (continued)

When using this product

- do not exceed recommended dosage
- marked drowsiness may occur
- excitability may occur, especially in children
- sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occurs
- symptoms do not improve within 7 days or occur with a fever
- cough persists for more than 7 days, comes back, or occurs with a fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not give more than 6 doses in 24 hours unless directed by a doctor
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- shake well before using

Age	Dose
children 6 to under 12 years of age	10 mL every 4 hours
children 4 to under 6 years of age	do not use unless directed by a doctor
children under 4 years of age	do not use

Other information

- each 5 mL contains: sodium 3 mg
- store between 20-25°C (68-77°F). Do not refrigerate.
- protect from light

Inactive ingredients acesulfame potassium, citric acid, EDTA disodium, FD&C blue #1, FD&C red #40, flavor, maltitol, propylene glycol, purified water, sodium benzoate, sodium citrate

Questions or comments?

Call 1-877-753-3835 Monday-Friday 9AM-5PM EST

*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Children's Triaminic® Day Time & Night Time Cold & Cough.

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

DAY & NIGHT COMBO PACK



Compare to the active ingredients in Children's Triaminic® Day Time Cold & Cough*

Daytime

Children's Cold & Cough

DEXTROMETHORPHAN HBr
Cough suppressant
PHENYLEPHRINE HCl
Nasal decongestant

Relieves:
• Cough
• Stuffy nose
Alcohol 0.14%
For Ages 4 to 11 Years



Cherry Flavor
Dosage cup included

Actual Bottle Size on Side Panel

4 FL OZ (118 mL) + 4 FL OZ (118 mL)
TOTAL 8 FL OZ (236 mL)



Compare to the active ingredients in Children's Triaminic® Night Time Cold & Cough*

NDC 69042-422-08

Children's Nighttime Cold & Cough

DIPHENHYDRAMINE HCl
Antihistamine; Cough suppressant
PHENYLEPHRINE HCl
Nasal decongestant

Relieves:
• Cough
• Runny, stuffy nose
• Itchy throat
Alcohol free
For Ages 6 to 11 Years



Grape Flavor
Dosage cup included

Actual Bottle Size on Side Panel

Package Contains Two Bottles

Actual Size

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.
KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2019 CVS/pharmacy
CVS.com® 1-800-SHOP CVS V-13114



PLD-B319E
PLD-B320E
FC005008

Lot No.:
Exp. Date:

